

# EndoClot® Submucosal Injection Solution (SIS)

# **Quick Reference Guide**

CAUTION: This guide is only a summary of the instructions for use for the EndoClot® Submucosal Injection Solution. Be sure to follow the instruction manual included in the device packaging.

#### **Product inspection and components**

- · Inspect package for damage
- · Peel off cover



### Prepare the agent



1. Place vial in holder.



2. Inject 15ml (SIS-15) /30ml (SIS-30) of sterile 0.9% saline into vial.



3 Mix by shaking vial.

# EndoClot® Submucosal Injection Solution (SIS)

# **Quick Reference Guide**

CAUTION: This guide is only a summary of the instructions for use for the EndoClot® Submucosal Injection Solution. Be sure to follow the instruction manual included in the device packaging.

#### Prepare the agent (continued)



4. Attach needle to SIS syringe.

#### Draw the agent into syringe



**5.** Unscrew locking cap to free syringe plunger.

#### Draw the agent into syringe (continued)



6. Pull the syringe plunger to draw agent.



7. Remove needle from syringe, then purge any air in the syringe.





8. Re-screw the locking cap.

# EndoClot® Submucosal Injection Solution (SIS)

# **Quick Reference Guide**

CAUTION: This guide is only a summary of the instructions for use for the EndoClot® Submucosal Injection Solution. Be sure to follow the instruction manual included in the device packaging.

#### Connect injection needle





9. Connect to needle and tighten the Luer-lock.





10. Purge air from needle by winding plunger.

#### **Deploy**



**11** Pass the injection needle down the scope channel.

#### Injection





12. Wind plunger to inject agent as appropriate.

EndoClot Plus Co., Ltd

Olympus is a registered trademark of Olympus Corporation, Olympus America Inc., and/or their affiliates. | Medical devices listed may not be available for sale in all countries.

