EndoClot[®] Submucosal Injection Solution (SIS) Quick Reference Guide

CAUTION: This guide is only a summary of the instructions for use for the EndoClot[®] Submucosal Injection Solution. Be sure to follow the instruction manual included in the device packaging.

Product inspection and components

- Inspect package for damage
- Peel off cover



Spiral dispensing syringe

Vial containing 0.5g (SIS-15)/1g (SIS-30) AMP® powder

Plastic tray with holder for vial

Prepare the agent



1 Place vial in holder.



2 Inject 15ml (SIS-15) /30ml (SIS-30) of sterile 0.9% saline into vial.



3 Mix by shaking vial.

EndoClot[®] Submucosal Injection Solution (SIS)

Quick Reference Guide

CAUTION: This guide is only a summary of the instructions for use for the EndoClot[®] Submucosal Injection Solution. Be sure to follow the instruction manual included in the device packaging.

Prepare the agent (continued)



Draw the agent into syringe



4 Attach needle to SIS syringe.

Draw the agent into syringe (continued)

5 Unscrew locking cap to free syringe plunger.

<image><image><image><image>

EndoClot[®] Submucosal Injection Solution (SIS)

Quick Reference Guide


CAUTION: This guide is only a summary of the instructions for use for the EndoClot[®] Submucosal Injection Solution. Be sure to follow the instruction manual included in the device packaging.

Connect injection needle



9 Connect to needle and tighten the Luer-lock.






11 Pass the injection needle down the scope channel.



Injection



M00511EN · 03/23 · OKM

As medical knowledge is constantly growing, technical modifications or changes of the product design, product specifications, accessories and service offerings may be required.



OLYMPUS EUROPA SE & CO. KG Postbox 10 49 08, 20034 Hamburg, Germany Wendenstrasse 20, 20097 Hamburg, Germany Phone: +49 40 23773-0, Fax: +49 40 233765 www.olympus-europa.com